

SIT-LP-08/05
May, 2008

On Wess-Zumino gauge

KAZUNARI SHIMA ^{*} and MOTOMU TSUDA [†]

*Laboratory of Physics, Saitama Institute of Technology
Fukaya, Saitama 369-0293, Japan*

Abstract

In the relation between the linear (L) supersymmetry (SUSY) representation and the nonlinear (NL) SUSY representation we discuss the role of the Wess-Zumino gauge. We show in two dimensional spacetime that a spontaneously broken LSUSY theory with mass and Yukawa interaction terms for a minimal off-shell vector supermultiplet is obtained from a general superfield without imposing any special gauge conditions in $N = 2$ NL/L SUSY relation.

PACS: 11.30.Pb, 12.60.Jv, 12.60.Rc, 12.10.-g

Keywords: supersymmetry, superfield, Nambu-Goldstone fermion, nonlinear/linear SUSY relation, composite unified theory

^{*}e-mail: shima@sit.ac.jp

[†]e-mail: tsuda@sit.ac.jp

Superfields on superspace provide a method to formulate SUSY field theories systematically [1, 2]. The Wess-Zumino (WZ) gauge is used conveniently to eliminate redundant (gauge) degrees of freedom (d.o.f.) from general superfields for vector supermultiplets. Linear (L) SUSY actions for minimal off-shell vector supermultiplets (in interacting theories) are constructed by using superfields in the WZ gauge.

On the other hand, various LSUSY theories are related (equivalent) to a NL SUSY model [3] through the linearization of NLSUSY both for the free case [4]-[8] and for the interacting (SUSY-QED) case [9, 10], which are essential to study the low energy physics and cosmology of NLSUSY general relativity (GR) [11] in the SGM scenario [12]. In the NL/L SUSY relation, all component fields for the vector supermultiplets including the redundant ones of the general superfields are consistently expressed as composites of Nambu-Goldstone (NG) fermion [7, 13] (called *superon* in the SGM scenario), whose expressions are called *SUSY invariant relations*. This means the WZ gauge condition used in LSUSY theories gives apparently the artificial (gauge dependent) restrictions on the composite states of NG fermions, which looks unnatural so far provided the composites are regarded as some eigen state of spacetime symmetry. Therefore, it is worthwhile studying the general properties of the superfield formulation which does not rely on the special gauge condition in viewpoints of NL/L SUSY relation.

In this letter, we focus on the interacting SUSY theory for the $N = 2$ vector supermultiplet in two dimensional spacetime ($d = 2$) for simplicity. Note that $N = 1$ vector supermultiplet is unphysical [14] in the SGM scenario. We show that by using the $N = 2$ NL/L SUSY relation constructed systematically in the previous paper [13] the interacting SUSY theory for the minimal off-shell vector supermultiplet and for the general vector supermultiplet as well accompanying the spontaneous SUSY breaking are obtained from the general superfield without imposing any special gauge conditions.

The $N = 2$ general superfield on superspace coordinates (x^a, θ^i) for the $N = 2$ LSUSY vector supermultiplet (in $d = 2$) is given by [15, 16]

$$\begin{aligned} \mathcal{V}(x, \theta^i) = & C(x) + \bar{\theta}^i \Lambda^i(x) + \frac{1}{2} \bar{\theta}^i \theta^j M^{ij}(x) - \frac{1}{2} \bar{\theta}^i \theta^i M^{jj}(x) + \frac{1}{4} \epsilon^{ij} \bar{\theta}^i \gamma_5 \theta^j \phi(x) \\ & - \frac{i}{4} \epsilon^{ij} \bar{\theta}^i \gamma_a \theta^j v^a(x) - \frac{1}{2} \bar{\theta}^i \theta^i \bar{\theta}^j \lambda^j(x) - \frac{1}{8} \bar{\theta}^i \theta^i \bar{\theta}^j \theta^j D(x), \end{aligned} \quad (1)$$

where $M^{ij} = M^{(ij)}$ ($= \frac{1}{2}(M^{ij} + M^{ji})$) and $M^{ii} = \delta^{ij} M^{ij}$. LSUSY transformations for the (general) component fields $(C, \Lambda^i, M^{ij}, \dots)$ with constant (Majorana) spinor parameters ζ^i are deduced from the following supertranslation of the superfield in the superspace (x^a, θ^i) ,

$$\delta_\zeta \mathcal{V}(x, \theta^i) = \bar{\zeta}^i Q^i \mathcal{V}(x, \theta^i) \quad (2)$$

with supercharges $Q^i = \frac{\partial}{\partial \theta^i} + i \not{\partial} \theta^i$. Gauge transformations for $(C, \Lambda^i, M^{ij}, \dots)$ are also defined as follows by means of [16]

$$\delta_g \mathcal{V}(x, \theta^i) = \Phi^1(x, \theta^i) + \Phi^2(x, \theta^i), \quad (3)$$

where Φ^i ($i = 1, 2$) are two scalar superfield parameters for generalized gauge parameters (for a $N = 2$ matter scalar supermultiplet (for example, see [6])).

As in the case for $d = 4$, $N = 1$ SUSY [1], gauge invariant quantities for $N = 2$ denoted tentatively as $(A_0, \phi_0, F_{0ab}, \lambda_0, D_0)$ can be easily constructed by the explicit component form of the gauge transformation (3) as follows;

$$(A_0, \phi_0, F_{0ab}, \lambda_0^i, D_0) \equiv (M^{ii}, \phi, F_{ab}, \lambda^i + i \not{\partial} \Lambda^i, D + \square C), \quad (4)$$

where $F_{0ab} = \partial_a v_{0b} - \partial_b v_{0a}$, $F_{ab} = \partial_a v_b - \partial_b v_a$ and $v_{0a} = v_a$ transforms as an Abelian gauge field. The familiar LSUSY transformations are calculated directly from Eq.(2). And for the minimal off-shell vector supermultiplet $(A_0, \phi_0, v_{0a}, \lambda_0^i, D_0)$ [9], we obtain

$$\begin{aligned} \delta_\zeta A_0 &= \bar{\zeta}^i \lambda_0^i, \\ \delta_\zeta \phi_0 &= -\epsilon^{ij} \bar{\zeta}^i \gamma_5 \lambda_0^j, \\ \delta_\zeta v_{0a} &= -i \epsilon^{ij} \bar{\zeta}^i \gamma_a \lambda_0^j + \partial_a W(\Lambda^i), \\ \delta_\zeta \lambda_0^i &= (D_0 - i \not{\partial} A_0) \zeta^i - i \epsilon^{ij} \gamma_5 \not{\partial} \phi_0 \zeta^j + \frac{1}{2} \epsilon^{ab} \epsilon^{ij} F_{0ab} \gamma_5 \zeta^j, \\ \delta_\zeta D_0 &= -i \bar{\zeta}^i \not{\partial} \lambda_0^i, \end{aligned} \quad (5)$$

where remarkably the redundant components in Eq.(1), $(C, \Lambda^i, M^{12}, M^{11} - M^{22})$, do not appear apparently in the minimal off-shell vector multiplet $\delta_\zeta(A_0, \phi_0, v_{0a}, \lambda_0^i, D_0)$ disregarding the gauge choice except for the $U(1)$ gauge parameter $W(\Lambda^i) = -2\epsilon^{ij} \bar{\zeta}^i \Lambda^j$ composed of the fermionic d.o.f.. That is, the gauge invariant sector and the redundant sector are mixed by only the gauge transformations with the fermionic parameters. Therefore the redundant sector plays important roles for constructing SUSY QED.

The LSUSY transformations (5) satisfy the closed commutator algebra,

$$[\delta_{\zeta_1}, \delta_{\zeta_2}] = \delta_P(\Xi^a), \quad (6)$$

where $\delta_P(\Xi^a)$ means a translation with a parameter $\Xi^a = 2i \bar{\zeta}_1^i \gamma^a \zeta_2^i$. Interestingly, in contrast to the WZ gauge, the commutator algebra for v_{0a} does not contain the

$U(1)$ gauge transformation term due to

$$\delta_{\zeta_1} W_{\zeta_2}(\Lambda^i) - \delta_{\zeta_2} W_{\zeta_1}(\Lambda^i) = -2(\epsilon^{ij} \bar{\zeta}_1^i \zeta_2^j A_0 + \bar{\zeta}_1^i \gamma_5 \zeta_2^i \phi_0 - i \bar{\zeta}_1^i \gamma^a \zeta_2^i v_{0a}) \quad (7)$$

and the consequent cancellations.

Now we see the above general arguments in the viewpoints of NL/L SUSY relation. Let us introduce the SUSY invariant relations between $(C, \Lambda^i, M^{ij}, \dots)$ of $N = 2$ LSUSY and (Majorana) NG fermions ψ^i of $N = 2$ NLSUSY [13] in $d = 2$, in which each component field is expanded in terms of ψ^i with respect to a constant κ whose dimension is $(\text{mass})^{-1}$ ($\kappa^{-2} \sim \frac{\Lambda}{G}$ in NLSUSY GR [11]) as

$$(C, \Lambda^i, M^{ij}, \dots) \sim \xi \kappa^{n-1} (\psi^i)^n |w| \quad (n = 4, 3, \dots, 0), \quad (8)$$

where ξ is an arbitrary dimensionless constant, $(\psi^i)^4 = \bar{\psi}^i \psi^i \bar{\psi}^j \psi^j$, $(\psi^i)^3 = \psi^i \bar{\psi}^j \psi^j$ and $(\psi^i)^2 = \bar{\psi}^i \psi^j$, $\epsilon^{ij} \bar{\psi}^i \gamma_5 \psi^j$, $\epsilon^{ij} \bar{\psi}^i \gamma^a \psi^j$, which are very promising features for SGM scenario [12]. In Eq.(8), $|w|$ is the determinant introduced in [3] describing the dynamics of NG fermions of NLSUSY, i.e. for the $d = 2$, $N = 2$ ($N > 2$, as well) NLSUSY case,

$$|w| = \det(w^a_b) = \det(\delta^a_b + t^a_b), \quad t^a_b = -i\kappa^2 \bar{\psi}^i \gamma^a \partial_b \psi^i, \quad (9)$$

expanded in terms of t^a_b or ψ^i as

$$\begin{aligned} |w| &= 1 + t^a_a + \frac{1}{2!} (t^a_a t^b_b - t^a_b t^b_a) \\ &= 1 - i\kappa^2 \bar{\psi}^i \not{\partial} \psi^i - \frac{1}{2} \kappa^4 (\bar{\psi}^i \not{\partial} \psi^i \bar{\psi}^j \not{\partial} \psi^j - \bar{\psi}^i \gamma^a \partial_b \psi^i \bar{\psi}^j \gamma^b \partial_a \psi^j) \\ &= 1 - i\kappa^2 \bar{\psi}^i \not{\partial} \psi^i - \frac{1}{2} \kappa^4 \epsilon^{ab} (\bar{\psi}^i \psi^j \partial_a \bar{\psi}^i \gamma_5 \partial_b \psi^j + \bar{\psi}^i \gamma_5 \psi^j \partial_a \bar{\psi}^i \partial_b \psi^j). \end{aligned} \quad (10)$$

Note that the NG fermion fields ψ^i transform nonlinearly as

$$\delta_\zeta \psi^i = \frac{1}{\kappa} \zeta^i - i\kappa \bar{\zeta}^j \gamma^a \psi^j \partial_a \psi^i, \quad (11)$$

which also satisfy the commutator algebra (6).

Explicit forms of the SUSY invariant relations in all orders of ψ^i of Eq.(8) are given as follows;

$$C = -\frac{1}{8} \xi \kappa^3 \bar{\psi}^i \psi^i \bar{\psi}^j \psi^j,$$

$$\begin{aligned}
\Lambda^i &= -\frac{1}{2}\xi\kappa^2\psi^i\bar{\psi}^j\psi^j(1-i\kappa^2\bar{\psi}^k\bar{\partial}\psi^k), \\
M^{ij} &= \frac{1}{2}\xi\kappa\bar{\psi}^i\psi^j\left(1-i\kappa^2\bar{\psi}^k\bar{\partial}\psi^k-\frac{1}{2}\kappa^4\epsilon^{ab}\bar{\psi}^k\psi^l\partial_a\bar{\psi}^k\gamma_5\partial_b\psi^l\right), \\
\phi &= -\frac{1}{2}\xi\kappa\epsilon^{ij}\bar{\psi}^i\gamma_5\psi^j\left(1-i\kappa^2\bar{\psi}^k\bar{\partial}\psi^k-\frac{1}{2}\kappa^4\epsilon^{ab}\bar{\psi}^k\gamma_5\psi^l\partial_a\bar{\psi}^k\partial_b\psi^l\right), \\
v^a &= -\frac{i}{2}\xi\kappa\epsilon^{ij}\bar{\psi}^i\gamma^a\psi^j(1-i\kappa^2\bar{\psi}^k\bar{\partial}\psi^k), \\
\lambda^i &= \xi\psi^i|w|, \\
D &= \frac{\xi}{\kappa}|w|.
\end{aligned} \tag{12}$$

The SUSY invariant relations (12) are systematically obtained [13] by considering the general superfield,

$$\tilde{\mathcal{V}}(x, \theta^i) = \mathcal{V}(x', \theta'^i), \tag{13}$$

on the specific supertranslations (x'^a, θ'^i) [4, 6] depending on ψ^i ,

$$\begin{aligned}
x'^a &= x^a + i\kappa\bar{\theta}^i\gamma^a\psi^i, \\
\theta'^i &= \theta^i - \kappa\psi^i.
\end{aligned} \tag{14}$$

Expanding the superfield (13) in the power series of θ^i , i.e.

$$\begin{aligned}
\tilde{\mathcal{V}}(x, \theta^i) &= \tilde{C}(x) + \bar{\theta}^i\tilde{\Lambda}^i(x) + \frac{1}{2}\bar{\theta}^i\theta^j\tilde{M}^{ij}(x) - \frac{1}{2}\bar{\theta}^i\theta^i\tilde{M}^{jj}(x) + \frac{1}{4}\epsilon^{ij}\bar{\theta}^i\gamma_5\theta^j\tilde{\phi}(x) \\
&\quad - \frac{i}{4}\epsilon^{ij}\bar{\theta}^i\gamma_a\theta^j\tilde{v}^a(x) - \frac{1}{2}\bar{\theta}^i\theta^i\bar{\theta}^j\tilde{\lambda}^j(x) - \frac{1}{8}\bar{\theta}^i\theta^i\bar{\theta}^j\theta^j\tilde{D}(x),
\end{aligned} \tag{15}$$

and $\mathcal{V}(x', \theta'^i)$ as well and imposing SUSY invariant constraints,

$$\tilde{C} = \tilde{\Lambda}^i = \tilde{M}^{ij} = \tilde{\phi} = \tilde{v}^a = \tilde{\lambda}^i = 0, \quad \tilde{D} = \frac{\xi}{\kappa}, \tag{16}$$

give the SUSY invariant relations (12). Note that the SUSY invariant constraints (16) do not restrict the NG fermion-composite structure of LSUSY multiplet fields.

The SUSY invariant relations for the component fields $(A_0, \phi_0, v_{0a}, \lambda_0, D_0)$ constituting the minimal off-shell vector supermultiplet are constructed from Eq.(12). And they coincide with the results obtained heuristically in Ref.[9], which induce the SUSY transformations (5) and the commutator algebra (6) under the NLSUSY transformations (11).

Now the problem is actions constructed from the general superfield (1) in NL/L SUSY relation. The $N = 2$ LSUSY free (kinetic) action with a Fayet-Iliopoulos (FI) term is written as

$$\begin{aligned}
S_{\mathcal{V}\text{free}} &= \int d^2x \left\{ \int d^2\theta^i \frac{1}{32} (\overline{D^i \mathcal{W}^{jk}} D^i \mathcal{W}^{jk} + \overline{D^i \mathcal{W}_5^{jk}} D^i \mathcal{W}_5^{jk}) + \int d^4\theta^i \frac{\xi}{2\kappa} \mathcal{V} \right\}_{\theta^i=0} \\
&= \int d^2x \left\{ -\frac{1}{4} (F_{0ab})^2 + \frac{i}{2} \bar{\lambda}_0^i \not{\partial} \lambda_0^i + \frac{1}{2} (\partial_a A_0)^2 + \frac{1}{2} (\partial_a \phi_0)^2 + \frac{1}{2} D_0^2 \right. \\
&\quad \left. - \frac{\xi}{\kappa} (D_0 - \square C) \right\} \\
&= S_{\mathcal{V}\text{free}}^0 + \int d^2x \frac{\xi}{\kappa} \square C,
\end{aligned} \tag{17}$$

where $\mathcal{W}^{ij}$ and $\mathcal{W}_5^{ij}$ are scalar and pseudo scalar superfields defined by

$$\mathcal{W}^{ij} = \bar{D}^i D^j \mathcal{V}, \quad \mathcal{W}_5^{ij} = \bar{D}^i \gamma_5 D^j \mathcal{V} \tag{18}$$

with $D^i = \frac{\partial}{\partial \theta^i} - i \not{\partial} \theta^i$. Obviously, the free action (17) is expressed by only the fields $(A_0, \phi_0, v_{0a}, \lambda_0, D_0)$ except the surface term without choosing the specific gauge. By changing the integration variables in Eq.(17) from (x, θ^i) to (x', θ'^i) under the SUSY invariant constraints (16) [13] or by directly substituting the SUSY invariant relations (12) into Eq.(17) [9], $S_{\mathcal{V}\text{free}}$ exactly reduces to the $N = 2$ NLSUSY action, i.e.

$$S_{\mathcal{V}\text{free}}(\psi^i) = S_{N=2\text{NLSUSY}}, \tag{19}$$

when $\xi^2 = 1$, where $S_{N=2\text{NLSUSY}}$ is

$$S_{N=2\text{NLSUSY}} = -\frac{1}{2\kappa^2} \int d^2x |w|, \tag{20}$$

which is invariant (becomes a surface term) under the NLSUSY transformations (11).

Next we focus on mass and Yukawa interaction terms constructed from Eq.(1). We define those terms by the quadratic and the cubic terms of $\mathcal{W}^{ij}$ and $\mathcal{W}_5^{ij}$ as follows;

$$\begin{aligned}
S_{\mathcal{V}m} &= \frac{1}{16} \int d^2x m \left[\int d^2\theta^i 3\{(\mathcal{W}^{jk})^2 + (\mathcal{W}_5^{jk})^2\} \right. \\
&\quad \left. + \int d\bar{\theta}^i d\theta^j 2(\mathcal{W}^{ik} \mathcal{W}^{jk} + \mathcal{W}_5^{ik} \mathcal{W}_5^{jk}) \right]_{\theta^i=0},
\end{aligned} \tag{21}$$

$$\begin{aligned}
S_{\mathcal{V}f} = & \frac{1}{8} \int d^2x \, f \left[\int d^2\theta^i \, \mathcal{W}^{jk} (\mathcal{W}^{jl} \mathcal{W}^{kl} + \mathcal{W}_5^{jl} \mathcal{W}_5^{kl}) \right. \\
& + \int d\bar{\theta}^i d\theta^j \, 2\{ \mathcal{W}^{ij} (\mathcal{W}^{kl} \mathcal{W}^{kl} + \mathcal{W}_5^{kl} \mathcal{W}_5^{kl}) \\
& \left. + \mathcal{W}^{ik} (\mathcal{W}^{jl} \mathcal{W}^{kl} + \mathcal{W}_5^{jl} \mathcal{W}_5^{kl}) \} \right]_{\theta^i=0}, \tag{22}
\end{aligned}$$

where f is an arbitrary constant with the dimension (mass)¹. The actions (21) and (22) contain the redundant fields in $\mathcal{V}(x, \theta^i)$ but these actions give the $N = 2$ LSUSY mass and the Yukawa interaction terms for the minimal off-shell vector supermultiplet (by imposing the WZ gauge condition) [9]. In NL/L SUSY relation, it can be easily shown that the actions (21) and (22) vanish, i.e.

$$S_{\mathcal{V}m}(\psi^i) = 0, \quad S_{\mathcal{V}f}(\psi^i) = 0, \tag{23}$$

by changing the integration variables in Eqs.(21) and (22) from (x, θ^i) to (x', θ'^i) under the SUSY invariant constraints (16).

The results (23) mean the followings. The explicit (general) component form of Eqs.(21) and (22) become

$$\begin{aligned}
S_{\mathcal{V}m} = & \int d^2x \, m \left\{ -\frac{1}{2} (\bar{\lambda}_0^i \lambda_0^i - 2A_0 D_0 + \epsilon^{ab} \phi_0 F_{0ab}) - 2(2A_0 \square C - \partial_a \bar{\Lambda}^i \gamma^a \not{\partial} \Lambda^i) \right\} \\
= & S_{\mathcal{V}m}^0 - 2 \int d^2x \, m (2A_0 \square C - \partial_a \bar{\Lambda}^i \gamma^a \not{\partial} \Lambda^i), \tag{24}
\end{aligned}$$

$$\begin{aligned}
S_{\mathcal{V}f} = & \int d^2x \, f [(A_0 \bar{\lambda}_0^i \lambda_0^i + \epsilon^{ij} \phi_0 \bar{\lambda}_0^i \gamma_5 \lambda_0^j - A_0^2 D_0 + \phi_0^2 D_0 + \epsilon^{ab} A_0 \phi_0 F_{0ab}) \\
& + \{A_0 \bar{\lambda}^i \lambda^i - 2M^{ij} \bar{\lambda}^i \lambda^j - 2A_0^2 D + 4(M^{ij})^2 D + \dots\}] \\
= & S_{\mathcal{V}f}^0 + \int d^2x \, f \{A_0 \bar{\lambda}^i \lambda^i - 2M^{ij} \bar{\lambda}^i \lambda^j - 2A_0^2 D + 4(M^{ij})^2 D + \dots\}, \tag{25}
\end{aligned}$$

where $S_{\mathcal{V}m}^0$ and $S_{\mathcal{V}f}^0$ are the mass and the Yukawa interaction terms for the minimal off-shell vector supermultiplet, which are invariant under the LSUSY transformations (5), respectively. By directly substituting Eq.(12) into $S_{\mathcal{V}m}^0$ and $S_{\mathcal{V}f}^0$, the (nontrivial) vanishments of the actions,

$$S_{\mathcal{V}m}^0(\psi^i) = 0, \quad S_{\mathcal{V}f}^0(\psi^i) = 0, \tag{26}$$

are shown by means of miraculous cancellations among four NG fermion self- interaction terms [9]. Therefore, it is understood that the redundant terms in the actions (24) and (25) also vanish by means of the cancellations among terms, which can be confirmed easily.

The above arguments show that, under the SUSY invariant relations (12), the $N = 2$ LSUSY action with the mass and the Yukawa interaction terms for the minimal off-shell vector supermultiplet $(A_0, \phi_0, v_{0a}, \lambda_0, D_0)$ is obtained from the NLSUSY action (20) without imposing any special gauge conditions for the general superfield (1); namely, the (nontrivial) vanishment of the redundant (subsidiary) structure of the action in NL/L SUSY relation in addition to Eqs.(19), (23) and (26) leads to

$$\begin{aligned} S_{N=2\text{NLSUSY}} &= S_{\mathcal{V}\text{free}} + S_{\mathcal{V}m} + S_{\mathcal{V}f} \\ &= S_{\mathcal{V}\text{free}}^0 + S_{\mathcal{V}m}^0 + S_{\mathcal{V}f}^0 + [\text{surface term}]. \end{aligned} \quad (27)$$

To summarize, based on the general superfield (1), we have discussed the relation between the $N = 2$ NLSUSY model and the $N = 2$ LSUSY theory for the vector supermultiplet with the mass and the Yukawa interaction terms in $d = 2$. The SUSY invariant relations and the SUSY invariant constraints in NL/L SUSY relation are expressed on the most general supermultiplet of the vector superfield. The SUSY invariant relations (12), which are systematically obtained [13] from the SUSY invariant constraints (16), are compatible with the gauge invariances of the quantities (4) for the minimal off-shell vector supermultiplet [9]. As for the actions constructed from Eq.(1), in particular, the mass and the Yukawa interaction terms (21) and (22), we have shown that the redundant (subsidiary) structure of the actions vanishes in NL/L SUSY relation in addition to Eqs.(19), (23) and (26). This means Eq.(27), i.e. the $N = 2$ LSUSY action with the mass and the Yukawa interaction terms for the minimal off-shell vector supermultiplet emerges from the NLSUSY action (20) without imposing any special gauge conditions for the general superfield (1).

The same structure as the $N = 2$ NL/L SUSY relation is anticipated for higher N SUSY and in $d = 4$ as well. The similar arguments in the coupling of matter multiplet (SUSY QED) and in the SUSY QCD are also interesting problems in NL/L SUSY relation and are under the study.

References

- [1] J. Wess and J. Bagger, *Supersymmetry and Supergravity (Second Edition, Revised and Expanded)* (Princeton University Press, Princeton, New Jersey, 1992).
- [2] P. West, *Introduction to Supersymmetry and Supergravity (Extended Second Edition)* (World Scientific, Singapore, 1990).
- [3] D.V. Volkov and V.P. Akulov, *Phys. Lett.* **B46** (1973) 109.
- [4] E.A. Ivanov and A.A. Kapustnikov, Relation between linear and nonlinear realizations of supersymmetry, JINR Dubna Report No. E2-10765, 1977 (unpublished);
E.A. Ivanov and A.A. Kapustnikov, *J. Phys.* **A11** (1978) 2375.
- [5] M. Roček, *Phys. Rev. Lett.* **41** (1978) 451.
- [6] T. Uematsu and C.K. Zachos, *Nucl. Phys.* **B201** (1982) 250.
- [7] K. Shima, Y. Tanii and M. Tsuda, *Phys. Lett.* **B525** (2002) 183;
- [8] K. Shima, Y. Tanii and M. Tsuda, *Phys. Lett.* **B546** (2002) 162.
- [9] K. Shima and M. Tsuda, *Mod. Phys. Lett.* **A22** (2007) 1085.
- [10] K. Shima and M. Tsuda, *Mod. Phys. Lett.* **A22** (2007) 3027.
- [11] K. Shima, *European Phys. J.* **C7** (1999) 341;
K. Shima, *Phys. Lett.* **B501** (2001) 237.
- [12] K. Shima and M. Tsuda, *Phys. Lett.* **B507** (2001) 260;
K. Shima and M. Tsuda, *Class. Quant. Grav.* **19** (2002) 5101;
K. Shima and M. Tsuda, *PoS HEP2005* (2006) 011.
- [13] K. Shima and M. Tsuda, Systematics of linearization of $N = 2$ nonlinear SUSY in superfield formulation, arXiv:0804.0846 [hep-th].
- [14] K. Shima and M. Tsuda, *Phys. Lett.* **B645** (2007) 455;
K. Shima, M. Tsuda and W. Lang, *Phys. Lett.* **B659** (2008) 741.
- [15] P. Di Vecchia and S. Ferrara, *Nucl. Phys.* **B130** (1977) 93.
- [16] K. Shima and M. Tsuda, On $N = 2$ superfield for $N = 2$ vector supermultiplet in two dimensional spacetime, arXiv:0802.0338 [hep-th], *Mod. Phys. Lett.* **A** (2008), at press.